

DYNAMICAL GENERATION OF THE PRIMORDIAL MAGNETIC FIELD BY FERROMAGNETIC DOMAIN WALLS

Paolo Cea^{a,b} and Luigi Tedesco^b

^aDipartimento di Fisica and ^bINFN, Sezione di Bari
Via Amendola 173, I-70126 Bari, Italy*

Abstract

The spontaneous generation of uniform magnetic condensate in QED_3 gives rise to ferromagnetic domain walls at the electroweak phase transition. These ferromagnetic domain walls are characterized by vanishing effective surface energy density avoiding, thus, the domain wall problem. Moreover we find that the domain walls generate a magnetic field $B \simeq 10^{24} Gauss$ at the electroweak scale which account for the seed field in the so called dynamo mechanism for the cosmological primordial magnetic field. We find that the annihilation processes of walls with size $R \simeq 10^5 Km$ could release an energy of order $10^{52} erg$ indicating the invisible ferromagnetic walls as possible compact sources of Gamma Ray Bursts.

Recently there was a renewed interest in the phenomenon of the spontaneous generation of the magnetic condensate both at zero [1] and finite [2, 3] temperature in three-dimensional spinorial electrodynamics. The interest in three-dimensional QED is not merely academic. Indeed, it is well known that Dirac fermions with a Yukawa coupling with a scalar field develop zero mode solutions near a domain wall [4]. These zero modes behave like massless fermions in the two-dimensional space of the wall. It has been suggested that ferromagnetic domain walls could be relevant for the formation of the primordial magnetic field [5]. The aim of this paper is to discuss a dynamical mechanism which induces an uniform magnetic condensate localized on the wall. Moreover, as we shall discuss, we find a natural solution of the domain wall problem so that we feel that our proposal provides a viable and realistic mechanism to account for the origin of the primordial magnetic field.

The only elementary scalar fields known in high energy physics are given by the scalar sector of Standard Model. According to the Standard Model when the Universe temperature reaches the electroweak scale $T_c \simeq 10^2 Gev$, there is a phase transition where the scalar field acquires a vacuum expectation value. In order to avoid inessential compliances we shall consider a simplified model which, however, retains the essential aspects of the full electroweak theory. In our model, the scalar sector is given by a real scalar field with Lagrangian:

$$\mathcal{L}_\phi = \frac{1}{2}(\partial_\mu \phi)^2 - \frac{\lambda}{4}(\phi^2 - v^2)^2. \quad (1)$$

At the phase transition the vacuum expectation value of the scalar field assumes the values $\langle \phi \rangle = \pm v$. So that one may assume that there are regions with $\langle \phi \rangle = +v$ and $\langle \phi \rangle = -v$. It is easy to see that the classical equation of motion admits the solution describing the transition layer between two regions with different values of $\langle \phi \rangle$:

$$\phi(z) = v \tanh\left(\frac{z}{\Delta}\right), \quad (2)$$

where

$$\Delta = \frac{1}{v\sqrt{\frac{\lambda}{2}}} \quad (3)$$

is the thickness of the wall. As it is well known there is some additional energy associated with the wall proportional to the area. The surface energy

density is given by:

$$\sigma_\phi = \int dz \left[\frac{1}{2}(\vec{\nabla}\phi)^2 + \frac{\lambda}{4}(\phi^2 - v^2)^2 \right] = \frac{2\sqrt{2}}{3}v^3\sqrt{\lambda}. \quad (4)$$

Let us consider, now, a four dimensional massless Dirac fermion coupled to the scalar field through the Yukawa coupling:

$$\mathcal{L}_\psi = \bar{\psi}(x)i \not{\partial} \psi(x) - g_Y \bar{\psi}(x)\psi(x)\phi(x). \quad (5)$$

In presence of the kink Eq.(2) the Dirac equation

$$[i \not{\partial} - g_Y \phi(x)]\psi(x) = 0 \quad (6)$$

admits the zero energy solutions localized on the wall [4]. Using the following representation of the Dirac algebra:

$$\gamma^3 = i \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \gamma^\alpha = i \begin{pmatrix} \tilde{\gamma}^\alpha & 0 \\ 0 & -\tilde{\gamma}^\alpha \end{pmatrix}, \quad \alpha = 0, 1, 2 \quad (7)$$

with

$$\tilde{\gamma}^0 = \sigma_3, \quad \tilde{\gamma}^1 = i\sigma_1, \quad \tilde{\gamma}^2 = i\sigma_2, \quad (8)$$

we find for the zero modes:

$$\psi_0(x, y, z) = \frac{N}{\sqrt{2}} \omega(z) \begin{pmatrix} \rho(x, y) \\ \rho(x, y) \end{pmatrix} \quad (9)$$

$$\omega(z) = \exp \left\{ -g_Y \int_0^z \phi(z') dz' \right\} \quad (10)$$

where $\rho(x, y)$ is a Pauli spinor, and

$$N = \left[\int_{-\infty}^{+\infty} dz \omega^2(z) \right]^{-\frac{1}{2}} = \sqrt{\pi} \Delta \frac{\Gamma(g_Y \Delta)}{\Gamma(g_Y \Delta + \frac{1}{2})}. \quad (11)$$

It is easy to see that $\rho(x, y)$ satisfies the 2 + 1-dimensional massless Dirac equation with the Dirac algebra Eq. (8).

After the electroweak spontaneous symmetry breaking the unique long range gauge field is the familiar electromagnetic field A_μ . In our scheme the real

scalar field is the physical Higgs field, so that $A_\mu(x)$ does not couple to $\phi(x)$. Thus the relevant Lagrangian turns out to be:

$$\mathcal{L} = -\frac{1}{4}F_{\mu\nu}(x)F^{\mu\nu}(x) + \bar{\psi}(x)[i \not{\partial} - e \not{A}(x)]\psi(x) - g_Y \bar{\psi}(x)\psi(x)\phi(x). \quad (12)$$

We are interested in the case of an external magnetic field localized on the wall at $z = 0$ and directed along the z -direction:

$$A_\mu(\vec{x}) = \left(0, \frac{B}{2}y, -\frac{B}{2}x, 0\right). \quad (13)$$

Moreover we shall restrict our analysis to the low-lying modes localized on the wall. In these approximations it is straightforward to check that the Hamiltonian reduces to:

$$H = \int d^3x \frac{B^2}{2} + \int d^2x \rho^\dagger(x, y) H^{(2)} \rho(x, y) \quad (14)$$

where

$$H^{(2)} = \tilde{\alpha}^k \cdot (i\partial_k - eA_k) \quad (15)$$

and $\tilde{\alpha}^k = \tilde{\gamma}^0 \tilde{\gamma}^k$, $k = 1, 2$. If we take into account the finite width of the wall Δ we may assume that the magnetic field B is sizeable over a distance $z \simeq \Delta$. So that Eq.(14) can be rewritten as

$$H = A\Delta \frac{B^2}{2} + \int d^2x \rho^\dagger(x, y) H^2 \rho(x, y), \quad (16)$$

where A is the area of the wall. In the following we shall consider the surface energy density. Equations (15) and (16) show that the zero mode contribution to the energy is given by two-dimensional massless Dirac fermions. Note that the fermionic contribution to H does not depend on the width of the wall. In the case of two-dimensional massive Dirac fermions it is known since long time that it is energetically favorable the spontaneous generation of an uniform constant magnetic field [6]. Indeed the fermionic energy density is given by [6]:

$$\mathcal{E}_F(B) = \frac{|eB|^{\frac{3}{2}}}{2\pi} g(\xi), \quad \xi = \frac{|eB|}{m^2}. \quad (17)$$

Near the origin one finds:

$$\mathcal{E}_F(B) = -\frac{|eB|}{4\pi} |m| + \frac{1}{24\pi^2} \frac{(eB)^2}{|m|} \quad (18)$$

The negative linear term in Eq. (18) gives rise to the negative minimum in the vacuum energy density. It is remarkable that this phenomenon survives to the thermal corrections, even in the infinite temperature limit [3].

On the other hand it can be shown that in 2+1-dimensional QED in presence of an external omogeneous background magnetic field with massless Dirac fermions it is energetically favourable to generate a constant mass term. To see this we follow the Cornwall, Jackiw and Tomboulis formalism of the effective potential for composite operators [7]. If we restrict to the case of a constant Dirac mass term coupled with the local operator $\bar{\rho}(x)\rho(x)$, $\rho(x)$ being the three-dimensional Dirac field, then the relevant quantity turns out to be:

$$\Omega = \mathcal{E}(B, m) - \mathcal{E}(B, 0) \quad (19)$$

where $\mathcal{E}(B, m)$ and $\mathcal{E}(B, 0)$ are respectively the expectation value of the Hamiltonian Eq. (15) on the massive and massless ground state. After some calculations, which will be reported elsewhere, we find in the one-loop approximation:

$$\begin{aligned} \Omega(B, m) = & -\frac{|eB|}{2\pi} \sum_{n=1}^{\infty} \sqrt{2n|eB| + m^2} + m^2 \frac{|eB|}{2\pi} \sum_{n=1}^{\infty} \frac{1}{\sqrt{2n|eB| + m^2}} + \\ & -\frac{\sqrt{2}\zeta(\frac{3}{2})}{8\pi^2} |eB|^{\frac{3}{2}}. \end{aligned} \quad (20)$$

A dimensional analysis shows that $\Omega(B, m) = |m|^3 f(\xi)$, where we recall that $\xi = \frac{|eB|}{m^2}$. In Figure 1 we report $f(\xi)$ versus $|m|$ for fixed $|eB|$. We see clearly that the massive ground state lies below the massless ground state for any value of the mass. Moreover the minimum is obtained for infinite mass. The above analysis strongly suggests that in three dimensional QED with a massless Dirac fermion there is spontaneous generation of both a mass term for the fermion field and a uniform magnetic field. In this case, from Eqs. (16,18) we have for the surface energy density of the ground state ¹

$$\sigma_{QED} = \mathcal{E}_{QED}(B, m) = \Delta \frac{B^2}{2} - \frac{|eB|}{4\pi} |m| + \frac{1}{24\pi^2} \frac{|eB|^2}{|m|}. \quad (21)$$

¹Strictly speaking Eq. (21) is valid for $\frac{|eB|}{m^2} \ll 1$. As we shall see, our results show that indeed this is the physically relevant region.

Remarkably Eq. (21) display a negative minimum at

$$B^* = \frac{e|m|^2}{4\pi} \frac{1}{|m|\Delta + \frac{e^2}{12\pi}} \quad (22)$$

with

$$\sigma_{QED} = \mathcal{E}_{QED}(B^*, m) = -\frac{e^2}{32\pi^2} \frac{|m|^3}{|m|\Delta + \frac{e^2}{12\pi}}. \quad (23)$$

To summarize, we have found that in presence of the kink solution Eq. (2) of the Higgs field there are massless fermion modes localized on the wall of the kink. In the one-loop approximation the zero modes becoming massive generate a constant magnetic field on the wall lowering the surface energy density by the amount given by Eq.(23). Thus the total surface energy density of the wall is:

$$\sigma = \sigma_\phi + \sigma_{QED} = \frac{2\sqrt{2}}{3} v^3 \sqrt{\lambda} - \frac{e^2}{32\pi^2} \frac{|m|^3}{|m|\Delta + \frac{e^2}{12\pi}}. \quad (24)$$

where we recall that $\Delta = \frac{\sqrt{\lambda}}{v\sqrt{2}}$.

The negative surface energy density induced by the magnetic condensation increases with the fermion zero mode mass. However the stability of the whole system requires that $\sigma \geq 0$. As a consequences the strength of the magnetic field increases up to the value such that

$$\sigma = 0. \quad (25)$$

By using the standard values $v \simeq 250 \text{ Gev}$, $\alpha_{QED} = \frac{e^2}{4\pi} \simeq \frac{1}{137}$ and assuming $\lambda \simeq 0.5$ we find that Eq.(25) admits one and only one positive root for $|m|$:

$$|m| \simeq 1.7 \cdot 10^4 \text{ Gev}, \quad (26)$$

and

$$B^* \simeq 5 \cdot 10^4 \text{ Gev}^2 \simeq 5 \cdot 10^{24} \text{ Gauss}. \quad (27)$$

Note that $\frac{eB^*}{m^2} \ll 1$ so that the approximation implied in Eq. (21) is rather good.

It is worthwhile to stress the main results we obtained. The remarkable phenomenon of spontaneous generation of an uniform magnetic condensate

in $(2 + 1)$ -dimensional QED gives rise to ferromagnetic domain walls which are invisible as concern the gravitational interactions. Indeed, according to Eq.(25) the surface energy density of the ferromagnetic walls is zero, thus solving the gravitational domain wall problem. In addition our invisible domain walls generate a magnetic field at the electroweak scale whose strength Eq. (27) agrees remarkably well with the one needed for the initial seed field in the dynamo mechanism for generating the cosmological magnetic field [8]. Finally our proposal for the invisible ferromagnetic domain walls could be relevant for the problem of the Gamma Ray Bursts [9]. Indeed the collision of a kink with an anti-kink should proceed like the annihilation of a fermion-antifermion pair. In the annihilation processes the negative energy of the magnetic condensate becomes disposable for the production of gamma particles. The energy for these processes can be estimated from Eq. (23). Indeed, from Equations (26) and (27) we get

$$\sigma_{QED} \simeq -4.2 \cdot 10^7 Gev^3, \quad (28)$$


which corresponds to the huge amount of disposable energy per fm^2

$$E \simeq 1fm^2 \cdot |\sigma_{QED}| \simeq 10^9 Gev. \quad (29)$$

If we require that the released energy in such process is of order $10^{52}erg$, then according to Eq. (29) we find that the linear dimension of the wall should be about $10^5 Km$. So that we see that our proposal provides relatively compact sources to account for the Gamma Ray Burst.

References

- [1] Y. Hosotani, Phys. Lett. B **319** (1993) 332; Phys. Rev. D **51** (1995) 2022; D. Cangemi, E. D'Hoker, and G. Dunne, Phys. Rev. D **51** (1995) 2513; V.P. Gusynin, U.A. Miransky, and I.A. Shovkovy, Phys. Rev. Lett **73** (1994) 3499; **76** (1996) 1005; Nucl. Phys. B 462 (1996) 249; V. Zeitlin, Phys. Lett. B **352** (1995) 422; P. Cea, Phys. Rev. D **55** (1997) 7895.
- [2] A. Das and M. Hott, Phys. Rev. D **53** (1996) 2252; S. Kanemura and T. Matsushita, Phys. Rev. D **56** (1997) 1021.
- [3] P. Cea and L. Tedesco, Phys. Lett. B **425** (1998) 345.
- [4] R. Jackiw and C. Rebbi, Phys. Rev. D **13** (1976) 3398; A.T. Niemi and G.W. Semenoff, Phys. Rep. C **135** (1986) 100.
- [5] A. Iwazaki, Phys. Lett B **406** (1997) 304; Phys. Rev. D **56** 1997 2435.
- [6] P. Cea, Phys Rev. D **32** (1985) 2785; Phys. Rev. D **34** (1986) 3229.
- [7] J. Cornwall, R. Jackiw and E. Tomboulis, Phys. Rev. D **10** (1974) 2428.
- [8] For a review see O. Tornkvist, *Magnetic Fields from The electroweak Phase Transition*, hep-ph/9801286; Phys. Rev. D **58** (1998) and references therein.
- [9] For a recent review see T. Piran, *Gamma-Ray Burst and Related Phenomena*, astro-ph/9801001 and references therein.



The one-loop effective potential $\frac{\Omega(B,m)}{|m|^3}$ versus $|m|$ for $|eB| = 1$.